

# The Nutritional Characteristics of a Contemporary Diet Based Upon Paleolithic Food Groups

Loren Cordain, PhD<sup>1\*</sup>

<sup>1</sup>Department of Health and Exercise Science, Colorado State University,  
Fort Collins, Colorado

## ABSTRACT

### Purpose:

The intent of the present study was to examine the nutritional characteristics of a contemporary diet based upon Paleolithic food groups and to determine how these characteristics may impact the risk of chronic disease.

### Methods:

Nutritional software was employed to ascertain the macro and trace nutrient characteristics of a diet composed of commonly available modern foods, but devoid of processed foods, dairy products and cereal grains. The relative contribution of plant and animal foods to the experimental diet was based upon average values previously determined in 229 hunter gatherer societies.

### Results:

The analysis revealed that except for vitamin D, which would have been supplied by endogenous synthesis in hunter gatherers, it is entirely possible to consume a nutritionally balanced diet from contemporary foods that mimic

the food groups and types available during the Paleolithic. Despite the elimination of two major food groups, the trace nutrient density of the experimental diet remained exceptionally high. The macronutrient content of the experimental diet (38 % protein, 39 % fat, 23 % carbohydrate by energy) varied considerably from current western values.

### Conclusions:

Contemporary diets based upon Paleolithic food groups maintained both trace and macronutrient qualities known to reduce the risk of a variety of chronic diseases in western populations.

## INTRODUCTION:

There is a growing awareness among evolutionary biologists that humans like all species are genetically adapted to the environment of their ancestors—that is, to the environment that their ancestors survived in and the environment that consequently conditioned their genetic makeup.<sup>1-3</sup> At the same time, there is growing awareness that the profound changes in the environment (e.g. in diet and other lifestyle conditions) that began with the introduction of agriculture and animal husbandry 10,000 years ago occurred too recently on an evolutionary timescale for the human genome to adjust.<sup>1-3</sup> As a result of the mismatch between the contemporary human diet and our genetically determined physiology, many of the so-called diseases of civilization have emerged.<sup>4-8</sup> Previous studies have examined the dietary characteristics of humans living during the Paleolithic,<sup>6,9,10</sup> as well as of historically studied hunter-gatherer societies,<sup>11,12</sup> and their authors have suggested that

---

### \* Correspondence:

Loren Cordain, PhD

Department of Health and Exercise Science

Colorado State University

Fort Collins, CO 80526

Phone: 970-491-7436 Fax: 970-491-0445

E-mail: cordain@cahs.colostate.edu

the nutritional qualities of these diets may have therapeutic value in the treatment of chronic disease. Although it is no longer possible or practical for contemporary men and women in western, industrialized countries to adopt and follow the exact dietary patterns of humans living during the Paleolithic, it is certainly possible to emulate the essential characteristics of historically studied hunter-gatherer diets with common foods and food groups available in all supermarkets.

The intent of this study was to examine the nutritional qualities of a contemporary diet based upon Paleolithic food groups and to characterize how these qualities may impact health and well being.

## METHODOLOGY

### Formulation of a Contemporary Diet Based Upon Paleolithic Food Groups

In the United States and other western nations, foods generally are organized into one of five food groups: 1) bread, cereal, rice and pasta group, 2) fruit group, 3) vegetable group, 4) milk, yogurt and cheese group, and 5) meat, poultry, fish, dry beans, eggs & nuts group.<sup>13</sup> The formulation of a contemporary diet based upon Paleolithic foods necessarily excludes two of these major groups (grains and dairy) because these foods were rarely or never consumed by contemporary or Paleolithic hunter-gatherers.<sup>9,11,14-15</sup> Additionally, within food group #5, dry beans and legumes were not included in the analysis because, like cereal grains, these foods did not become dietary staples until Neolithic times.<sup>16</sup> Finally, all modern processed foods containing mixtures of grains, refined sugars and oils, salt, and food additives were likewise excluded from the model because these food mixtures became part of the human dietary repertoire only following the Agricultural and Industrial Revolutions.<sup>9,11,14-15</sup> Consequently, the present model utilized only the following contemporary food types: fruits, vegetables, meats, poultry, fish and nuts/seeds. For each food type, only the most commonly consumed foods in the U.S. diet were incorporated into the model. These were then randomly arranged into three meals and snacks utilizing dishes that were not dissimilar from those normally found in traditional western diets. The example diet was then analyzed for macro and trace nutrients using nutritional software (Nutritionist 5, First Data Bank, San Bruno, CA).


The 20 most commonly consumed fruits, vegetables, and fish in the United States were employed in the random meal selections (Table 1).<sup>17</sup> For the 20 most commonly consumed vegetable foods in the United States, two foods (potatoes and corn) were excluded from the model because corn is a cereal grain, and potatoes maintain nutrient properties (high glycemic and insulin responses)<sup>18</sup> uncharacteristic of traditional hunter-gatherer plant foods.<sup>19</sup> Consequently, the remaining 18 vegetable foods in Table 1 represent the food choices available in the model.

For the meat food group, the four most commonly consumed meats in the United States (beef, chicken, pork and turkey)<sup>20</sup> represented the meats of choice in the analysis. Only very lean cuts of meat (turkey and chicken breasts without skin, pork loin trimmed of fat, beef sirloin tip roast trimmed of fat) that averaged 20 % fat by energy—a mean value similar to that found in wild game meat<sup>21</sup>—were utilized in the model. For the nuts/seeds group, 10 nuts and seeds commonly consumed in the U.S. diet (almonds, walnuts, pecans, filberts, brazil nuts, pistachio nuts, macadamia nuts, coconut, sunflower seeds and pumpkin seeds) represented the available choices for this food type.

The primary consideration in the formulation of a “modern Paleolithic diet” is the relative contribution of each food group to total energy intake. Compiled ethnographic studies of 229 hunter-gatherer societies,<sup>11</sup> as well as 13 quantitative studies of hunter-gatherers<sup>12</sup> have demonstrated that animal foods contributed slightly more than half (55-65%) of the daily energy, whereas plant foods would have made up the remainder (35-45%) of the average daily caloric intake. Of the energy obtained from animal foods, historically-studied hunter-gatherers typically derived half of their energy from aquatic animals and the other half from terrestrial animals.<sup>11</sup> Animal food intake would have also been constrained by the physiologic protein ceiling, which has been shown to occur between 30 to 41% of total energy.<sup>11</sup>

In hunter-gatherer diets, the balance of total dietary energy (35-45%) derived from plant foods would have been quite erratic in how it would have been apportioned among the various plant food groups due to varying environmental and ecological considerations.<sup>11</sup> Hence, in the formulation of a modern diet based upon Paleolithic food groups, the plant food energy was arbitrarily divided equally among fruits, vegetables and nuts/seeds. Figure 1 displays the food type energy weightings assigned to the example “Modern Paleolithic” diet. Using these energy weightings for each of the five food types, the diet outlined in Table 2 was formulated.

**Figure 1.** Apportionment of daily energy to the five food types in a contemporary diet based upon Paleolithic food groups.



**Table 1.** Top 20 most common fruits, vegetables and fish sold in the United States.<sup>17</sup>

Top 20 Fruits	Top 20 Vegetables	Top 20 Fish
Banana	Potato	Shrimp
Apple	Iceberg lettuce	Cod
Watermelon	Tomato	Pollock
Orange	Onion	Catfish
Cantaloupe	Carrot	Scallop
Grape	Celery	Atlantic/Coho salmon
Grapefruit	Sweet corn	Flounder
Strawberry	Broccoli	Sole
Peach	Green cabbage	Oyster
Pear	Cucumber	Orange roughy
Nectarine	Bell pepper	Mackerel
Honeydew melon	Cauliflower	Ocean perch
Plum	Leaf lettuce	Rockfish
Avocado	Sweet potato	Whiting
Lemon	Mushroom	Clam
Pineapple	Green onion	Haddock
Tangerine	Green bean	Blue crab
Sweet cherry	Radish	Rainbow trout
Kiwi fruit	Summer squash	Halibut
Lime	Asparagus	Lobster

**Table 2.** Sample 1-day menu for a modern diet based upon Paleolithic food groups for females (25 yrs, 2200 kcal daily energy intake).

Breakfast	Food Quantity (g)	Energy (kcal)
Cantaloupe	276	97
Atlantic salmon (broiled)	333	605
<b>Lunch</b>		
Vegetable salad with walnuts		
Shredded Romaine lettuce	68	10
Sliced carrot	61	26
Sliced cucumber	78	10
Quartered tomatoes	246	52
Lemon juice dressing	31	8
Walnuts	11	70
Broiled lean pork loin	86	205
<b>Dinner</b>		
Vegetable avocado/almond salad		
Shredded mixed greens	112	16
Tomato	123	26
Avocado	85	150
Slivered almonds	45	260
Sliced red onion	29	11
Lemon juice dressing	31	8
Steamed broccoli	468	131
Lean beef sirloin tip roast	235	400
Dessert – Strawberries	130	39
<b>Snacks</b>		
Orange	66	30
Carrot sticks	81	35
Celery sticks	90	14

**Table 3.** Macronutrient and other dietary characteristics in a contemporary diet based on Paleolithic food groups for females (25 yrs, 2200 kcal daily energy intake).

Protein (g)	217
Protein (% energy)	38
Carbohydrate (g)	129
Carbohydrate (% energy)	23
Total sugars (g)	76.5
Fiber (g)	42.5
Fat (g)	100.3
Fat (% total energy)	39.0
Saturated fat (g)	18.0
Saturated fat (% total energy)	7.0
Monounsaturated fat (g)	44.3
Polyunsaturated fat (g)	26.7
Omega 3 fat (g)	9.6
Omega 6 fat (g)	14.2
Cholesterol (mg)	461
Sodium (mg)	726
Potassium (mg)	9062

## RESULTS

### Nutritional Characteristics of a Contemporary Paleolithic Diet

Table 3 presents the macronutrient intake and other qualities of the example diet analyzed from foods listed in Table 2. The macronutrient characteristics of the example diet, protein (38% energy), carbohydrate (23% energy), fat (39% energy) are similar to values demonstrated in historically studied hunter-gatherer societies but different from values (16% protein, 49% carbohydrate, 34% fat) in traditional western diets.<sup>11</sup> Despite its relatively low carbohydrate content (23% energy), the contemporary Paleolithic diet contained 42.5 g of plant fiber.

The contemporary Paleolithic diet contains more fat (39% energy) than average values (34% energy) found in western diets,<sup>11</sup> however this extra fat occurs entirely as a consequence of a greater intake of both polyunsaturated (PUFA) and monounsaturated (MUFA) fats. Although more than 50% of the energy in the contemporary Paleolithic diet is derived from animal foods, the saturated fat content (7.0% energy) falls within recommended healthful limits ( $\leq 10\%$  energy).<sup>13</sup> The contemporary Paleolithic diet is characterized by a high intake of total omega 3 (n3) fatty acids (9.6 g) and a relatively low intake of omega 6 (n6) fatty acids, which in turn yield a total dietary n6/n3 of 1.5 to 1. The cholesterol content of the contemporary Paleolithic diet is higher (461 mg) than recommended values (300 mg).<sup>13</sup> The contempo-

rary Paleolithic diet contains 12.5 times more potassium than sodium. Except for calcium, all trace nutrients occur in considerably greater quantities than the recommended daily allowances (RDAs) (Table 4).

## DISCUSSION

The results of this analysis demonstrate that it is entirely possible to consume a nutritionally balanced diet from commonly available contemporary foods that emulate the food types available to Paleolithic hunter gatherers. Despite the elimination of two major food groups (dairy and cereals), the trace nutrient density of the diet remains exceptionally high. Moreover, the diet maintains numerous nutritional characteristics that have been demonstrated to reduce the risk of a variety of chronic diseases.

### Potential Nutritional Shortcomings of the Contemporary Paleolithic Diet

#### Calcium

Table 4 shows that the calcium intake (691 mg) would be considerably lower than the RDA (1000 mg), while the protein intake (217 g) would be more than 4 times recommended values (50 g). Because increased dietary protein increases obligatory loss of urinary calcium, it has been suggested that a calcium (mg)/protein (g) ratio of  $\geq 20:1$  may protect against bone loss.<sup>22</sup> The calcium/protein ratio of the contemporary Paleolithic diet (3.2 :1) is considerably lower than that in the average U.S. diet (10.7:1)<sup>23</sup> and therefore might be expected to increase the risk for bone demineralization, osteoporosis, and osteopenia. However, analyses of the skeletons of ancestral humans living during the Paleolithic<sup>24,25</sup> as well as more recently studied hunter-gatherers<sup>26</sup> have shown these people maintained robust, fracture-resistant bones, free from signs and symptoms of osteoporosis despite consuming no dairy products. Their robust bones may be due in part to greater activity levels (bone loading)<sup>24</sup> and greater sunlight exposure (increased vitamin D synthesis, hence increased calcium absorption). However, more importantly it is likely that Paleolithic hunter gatherers would have been in positive calcium balance despite a relatively low calcium intake because the calciuretic effects of a high meat diet were countered by high fruit and vegetable intake.<sup>11,12</sup>

Ingestion of meat protein induces calciuresis because the oxidation of sulfur-containing amino acids presents a net acid load to the kidney, which in turn must buffer the acid load from base that ultimately is derived from calcium-containing bone mineral salts.<sup>27</sup> Previous studies have demonstrated that ingestion of an alkalizing agent prevented the calciurea which normally accompanies high protein diets,<sup>28</sup> and that when base is administered at a dose sufficient to neutralize endogenous acid production, calcium balance is improved, bone resorption is reduced, and bone formation is increased.<sup>29</sup> In western diets, meats,

**Table 4.** Trace nutrients in a modern diet based on Paleolithic food groups for females (25 yrs, 2200 kcal daily energy intake).

	Total	% RDA
Vitamin A (RE)	6386	798
Vitamin B <sub>1</sub> (mg)	3.4	309
Vitamin B <sub>2</sub> (mg)	4.2	355
Vitamin B <sub>3</sub> (mg)	60	428
Vitamin B <sub>6</sub> (mg)	6.7	515
Folate (µg)	891	223
Vitamin B <sub>12</sub> (µg)	17.6	733
Vitamin C (mg)	748	1247
Vitamin E (IU)	19.5	244
Calcium (mg)	691	69
Phosphorus (mg)	2546	364
Magnesium (mg)	643	207
Iron (mg)	24.3	162
Zinc (mg)	27.4	228

**Table 5.** The potential renal acid load (PRAL) in the example diet. Values for PRAL were adapted from Remer and Manz's database.<sup>30</sup> (+) values are acid-producing, (-) values are base-producing.

<u>Alkaline-Yielding Foods</u>	<u>Weight (g)</u>	<u>PRAL/100 g</u>	<u>Net PRAL</u>
Cantaloupe	276	-3.5	-9.7
Lettuce	68	-2.5	-1.7
Carrot	142	-4.9	-7.0
Cucumber	78	-0.8	-0.6
Tomatoes	331	-3.1	-10.3
Lemon juice	62	-2.5	-1.6
Salad greens	112	-2.5	-2.8
Avocado	85	-3.5	-3.0
Almonds	45	-2.8	-1.3
Onion	29	-1.5	-0.4
Broccoli	468	-1.2	-5.6
Strawberries	130	-2.2	-2.9
Orange	66	-2.7	-1.8
Celery	90	-5.2	-4.7
<b>Total</b>			<b>-53.2</b>
<u>Acid-Yielding Foods</u>			
Salmon	333	7.9	26.3
Pork	86	7.9	6.8
Beef	235	7.8	18.3
Walnuts	11	6.8	0.7
<b>Total</b>			<b>51.4</b>

cheeses, and cereal grains yield high potential renal acid loads<sup>30</sup> and hence may promote osteoporosis by producing a net metabolic acidosis.<sup>27</sup> In contrast, fruits and vegetables yield a net alkaline renal load,<sup>30</sup> and high fruit and vegetable diets have been shown to reduce urinary calcium excretion rates.<sup>31</sup> Accordingly, in hunter-gatherer populations consuming high protein diets, a concomitant consumption of high levels of fruits and vegetables may have countered the calciuretic effects of a high protein diet.

In the present model, the net renal ionic load was slightly alkaline with base producing foods (-53.2) outweighing acid producing foods (51.4) (Table 5). Consequently the high protein intake of the example diet would not have caused an increased calciuresis, and subjects consuming a similar diet likely would remain in calcium balance despite a calcium intake lower than the RDA.

### **Vitamin D**

The contemporary Paleolithic diet provides no dietary vitamin D. Except for the livers of certain marine mammals and fish, there are relatively few sources of vitamin D in the normal food supply. In most hunter-gatherers, vitamin D would have been obtained via the body's synthesis of this hormone from ultraviolet irradiation (sunlight exposure) of cholesterol in the skin. Only with the fortification of margarine and milk, beginning in the mid 20<sup>th</sup> century, has vitamin D been widely available in the food supply.

### **Cholesterol**

Table 3 shows that the cholesterol intake (461 mg) for the model diet is more than 50% higher than recommended values (300 mg).<sup>13</sup> However, it should be noted that dietary cholesterol has a relatively minor impact on serum cholesterol levels.<sup>32</sup> The recently developed Howell *et al.* equation<sup>33</sup> [ $\Delta$  serum cholesterol (mg/dL) = 1.918 x  $\Delta$  SFA - 0.900 x  $\Delta$  PUFA + 0.0222 x  $\Delta$  cholesterol; where SFA = % saturated fat energy, PUFA = % polyunsaturated fat energy, and cholesterol = dietary cholesterol (mg)] reveals that a reduction in dietary cholesterol from 461 mg (the value in the example diet) to 300 mg (recommended value) would only lower serum cholesterol levels by 3.5 mg/dL. Additionally, in the example diet the ratio of polyunsaturated fatty acids to saturated fatty acids (P/S) is 1.5.

Schonfeld and colleagues<sup>34</sup> have shown that when the P/S was = 0.8, the addition of 750 mg of dietary cholesterol did not elevate serum LDL cholesterol concentrations in healthy, normal men. Consequently, the high P/S in the contemporary Paleolithic diet likely would counter any elevations in serum cholesterol that potentially could have occurred from increased dietary cholesterol.

## **Potential Nutritional Benefits of the Modern Paleolithic Diet**

### **Dietary Protein**

Perhaps the most striking difference between the typi-

cal western diet and the current model diet lies in the much higher protein intake. Although a high protein ingestion can increase the rate of progression in renal dysfunction,<sup>35</sup> a recent clinical trial has demonstrated that a high protein diet (26% energy) had no adverse effects upon renal function in subjects with no pre-existing kidney disease.<sup>36</sup> Because protein has more than three times the thermic effect of either fat or carbohydrate<sup>37</sup> and because it has a greater satiety value than fat or carbohydrate,<sup>37,38</sup> increased dietary protein may represent an effective weight loss strategy for the overweight or obese. Recent clinical trials have demonstrated that calorie-restricted high protein diets are more effective than calorie-restricted high carbohydrate diets in eliciting weight loss in overweight subjects.<sup>39,40</sup>

There is an increasing body of evidence that suggests high protein diets may improve blood lipid profiles<sup>41-45</sup> and thereby lessen the risk for cardiovascular disease (CVD). Wolfe and colleagues have shown that the isocaloric substitution of protein (23% energy) for carbohydrate in moderately hypercholesterolemic subjects resulted in significant decreases in total, LDL and VLDL cholesterol, and TG while HDL cholesterol increased.<sup>43</sup> Similar blood lipid changes have been observed in type II diabetic patients in conjunction with improvements in glucose and insulin metabolism.<sup>41,42</sup> Further, high protein diets have been shown to improve metabolic control in type II diabetes patients.<sup>41,42,46</sup> In obese women, hypo-caloric high protein diets improved insulin sensitivity and prevented muscle loss, whereas hypo-caloric high carbohydrate diets worsened insulin sensitivity and caused reductions in the fat free mass.<sup>47</sup>

Epidemiological evidence supports the clinical data showing a cardiovascular protective effect of dietary protein. Increased protein intake has been shown to be inversely related to CVD in a cohort of 80,082 women.<sup>48</sup> Dietary protein is also inversely related to serum homocysteine concentration,<sup>49</sup> an independent risk factor for CVD. Meat eating populations have been shown to maintain lower plasma homocysteine concentrations than non-meat eaters.<sup>50,51</sup> In numerous population studies, summarized by Obarzanek *et al.*,<sup>52</sup> higher blood pressure was associated with lower intake of protein. Recently, a four-week dietary intervention of hypertensive subjects demonstrated that a high protein diet (25% energy) was effective in significantly lowering blood pressure.<sup>53</sup> Further, a number of population studies have established that stroke mortality is inversely related to protein intake.<sup>54,55</sup>

### **Dietary Carbohydrate and Fiber**

Table 3 reveals that the carbohydrate content (23% energy) of the example diet is considerably lower than average values (49% energy) in the U.S. diet,<sup>11</sup> or suggested healthful ranges (55-60% energy).<sup>13,56</sup> Although current advice to reduce the risk of CVD is, in general, to replace saturated fats with complex carbohydrate,<sup>13,56</sup> there is

mounting evidence to indicate that low fat, high carbohydrate diets may elicit undesirable blood lipid changes, including reductions in HDL cholesterol and apolipoprotein A-1, while concurrently elevating TG, VLDL cholesterol and small dense LDL cholesterol.<sup>57-60</sup> Collectively, these blood lipid changes are associated with an increased risk for CVD and other Syndrome X diseases.<sup>61</sup>

Table 6 shows both the glycemic index and glycemic load (glycemic index x carbohydrate content in food portion) in selected grain products, sugars/sweets, dairy foods, fruits, and vegetables.<sup>62</sup> High glycemic loads represent a nearly universal characteristic of the typical western diet because of a high reliance upon refined sugars and cereal grains. Added sugars represent 16.1% of the energy consumed in the average U.S. diet, whereas refined grain products comprise 85.3% of all the grains consumed in the U.S.<sup>23</sup> Table 6 reveals that dairy products maintain low glycemic indices and loads, but paradoxically these foods are highly insulinotropic with insulin indices similar to white bread.<sup>63</sup> Consequently, the elimination of refined sugars, grains and dairy products in the example diet produces a low-carbohydrate diet (23% energy) in which all of the carbohydrates are derived from fruits, vegetables, and seeds/nuts with their universally low glycemic loads. High glycemic load diets have been implicated in the development of obesity,<sup>64</sup> and observational studies suggest that foods with a high glycemic load increase the risk for type II diabetes<sup>65,66</sup> and CVD.<sup>67</sup>

The fiber content (42.5 g) of the example diet is considerably higher than values in the U.S. diet (15.1 g) and higher than recommended values (25-30 g).<sup>56</sup> Soluble fibers modestly reduce LDL and total cholesterol concentrations beyond those achieved by a diet low in saturated fat, and fiber, by slowing gastric emptying, may reduce appetite and help to control caloric intake.<sup>68</sup>

### Dietary Fat

The total fat content (39% energy) of the example diet is 30% higher than recommended intakes.<sup>13,56</sup> However, it should be noted that the overall dietary lipid profile is health-promoting and anti-atherogenic.

There is now substantial evidence to indicate that the absolute amount of dietary fat is less important in lowering blood lipid levels and reducing the risk for CVD than is the relative concentrations of specific dietary fatty acids.<sup>69-72</sup> Low (22% energy) and high (39% energy) fat diets which had identical (polyunsaturated/saturated) (n3/n6) and (monounsaturated/total fat) fatty acid ratios produced no significant differences in total or LDL cholesterol following a 50 day trial.<sup>72</sup> Hypercholesterolemic fatty acids include 12:0, 14:0, 16:0, and trans-9 18:1,<sup>73</sup> whereas monounsaturated (MUFA)<sup>70,74</sup> and polyunsaturated (PUFA)<sup>73</sup> fatty acids are hypocholesterolemic, and 18:0 is neutral.<sup>74</sup> Omega 3 PUFA have wide-ranging cardiovascular protective

**Table 6.** Glycemic indices and glycemic loads of various food groups. Glycemic load = (glycemic index x carbohydrate content in 10g portions). The glycemic reference is glucose with a glycemic index of 100. Data adapted from Foster-Powell *et al.*<sup>62</sup>

	Glycemic			Glycemic	
	Index	Load		Index	Load
<u>Grain Products</u>			<u>Vegetables</u>		
Rice krispie cereal	88	77.3	Parsnips	97	19.5
Cornflakes	84	72.7	Sweet potato	54	13.1
Rice cakes	82	66.9	Yam	51	11.5
Shredded wheat cereal	69	57.0	Carrots	71	7.2
Graham crackers	74	56.8	Beets	64	6.3
Cheerio cereal	74	54.2	Rutabaga	72	6.3
Rye crisp bread	65	53.4	<u>Fruits</u>		
Vanilla wafers	77	49.7	Banana	53	12.1
Corn chips	73	46.3	Pineapple	66	8.2
Wheat thins	67	41.9	Grapes	43	7.7
Granola bar	61	39.3	Kiwi fruit	52	7.4
Bagel	72	38.4	Apple	39	6.0
Doughnuts	76	37.8	Pear	36	5.4
White bread	70	34.7	Watermelon	72	5.2
All bran cereal	42	32.5	Orange	43	5.1
Whole wheat bread	69	31.8	<u>Dairy foods</u>		
<u>Sugar, sweets</u>			Ice cream	61	14.4
Jelly beans	80	74.5	Yogurt	33	6.5
Lifesavers	70	67.9	Skim milk	32	1.6
Table sugar (sucrose)	65	64.9	Whole Milk	27	1.3
Mars bar	68	42.2			

capacities including lowering of plasma VLDL cholesterol and triacylglycerol (TG) concentrations.<sup>69</sup> Consequently, it is entirely possible to consume relatively high fat diets that do not necessarily produce a plasma lipid profile that promotes CVD<sup>72,75</sup> given sufficient MUFA,<sup>70</sup> PUFA,<sup>60</sup> and an appropriate n6/n3 PUFA ratio<sup>69</sup> relative to the hypercholesterolemic fatty acids.

Although more than 50 % of the energy in the contemporary Paleolithic diet is derived from animal foods, the saturated fat content (7.0% energy) not only falls within recommended healthful limits ( $\leq 10$  % energy),<sup>13,56</sup> but also within limits ( $\leq 7$  %) for individuals with elevated LDL cholesterol concentrations or CVD.<sup>76</sup> The dominant fats in the example diet are cholesterol lowering MUFA (17.2 % energy) and PUFA (10.4 % energy). MUFA may also confer additional cardiovascular protective effects beyond lowering serum cholesterol by its ability to reduce LDL oxidizability, a key step in the atherosclerotic process.<sup>77</sup>

The example diet is rich in omega 3 fatty acids (9.6 g) compared to the average value (2.3 g) found in the U.S. diet.<sup>78</sup> Numerous studies have reported the beneficial effects of an increased omega 3 fatty acid intake in CVD patients.<sup>79-82</sup> A 20% reduction in overall mortality and a 45% reduction in sudden death after 3.5 years were reported in subjects with preexisting CVD when given 850 mg of omega 3 fatty acids (20:5n3 and 22:6n3) either with or without vitamin E.<sup>82</sup> Omega 3 fatty acids may operate to reduce CVD mortality via a number of mechanisms including reductions in serum VLDL and triacylglycerol concentrations, thrombotic tendencies, and the incidence of ventricular arrhythmias.<sup>66</sup>

### **Dietary Sodium and Potassium**

Because no processed foods or added salt are included in the example diet, the sodium intake (726 mg) is appreciably lower than average U.S. values (3,271 mg)<sup>23</sup> or recommended values (2,400 mg).<sup>56</sup> Further, since potassium-rich fruits and vegetables comprise 30% of the daily energy, the potassium content (9,062 mg) of the example diet is nearly 3.5 times greater than average values (2,620 mg) in the U.S. diet.<sup>23</sup> Diets rich in potassium and low in sodium have been repeatedly demonstrated to be therapeutic for a variety of chronic conditions including: hypertension, stroke, kidney stones, and osteoporosis.<sup>83,84</sup>

### **Trace Nutrients**

Table 4 demonstrates that, except for calcium, the example diet is exceedingly rich in the 14 vitamins and minerals most commonly deficient in the U.S. diet.<sup>23</sup> A meta-analysis investigating the relationship between CVD and serum homocysteine concentrations has demonstrated that as much as 10% of CVD risk was attributable to hyperhomocysteinemia.<sup>85</sup> The normal metabolism of homocysteine requires an adequate supply of folate, vitamin B<sub>6</sub>, vitamin B<sub>12</sub> and riboflavin. Lower serum folate concentrations and vitamin B<sub>6</sub> have been associated with increased CVD risk.<sup>86</sup> Because the fruits (15% energy) and vegetables (15% energy) in the example diet are rich sources of folate, the intake of this vitamin is quite high (891 µg or 223% RDA). Additionally, the fish (27.5% energy) and lean meats (27.5% energy) contained in the example diet are rich sources of vitamin B<sub>6</sub>, and along with the fruits, vegetables and seeds/nuts, combine to yield a high intake (6.7 mg or 515 % RDA) of this vitamin.

### **SUMMARY**

Despite a high reliance upon low fat animal foods (55% energy), the experimental diet would not have necessarily elicited unfavorable blood lipid profiles because of the hypolipidemic effects of high dietary protein (38 % energy) and the relatively low level of low glycemic index dietary carbohydrates (23%). Although total fat intake (39%

energy) would have been higher than that found in western diets, total saturated fat (7.0% energy) fell well within healthful limits (10% energy). Important qualitative differences in fat intake, including relatively high levels of MUFA and PUFA and a lower n6/n3 fatty acid ratio, also would have served to reduce the risk for CVD. Other characteristics of the example diet, including a high intake of antioxidants, fiber, vitamins, and phytochemicals along with a low salt intake would further deter the risk of CVD and other chronic diseases.

### **REFERENCES**

1. Williams GC, Nesse RM. The dawn of Darwinian medicine. *Quart Rev Biol.* 1991;66:1-22.
2. Wilson DR. Evolutionary epidemiology. *Acta Biotheoretica.* 1993;41:205-218.
3. Goldsmith MF. Ancestors may provide clinical answers, say 'Darwinian' medical evolutionists. *JAMA.* 1993;269:1477-1480.
4. Cohen MN. *Health and the Rise of Civilization.* London: Yale University Press; 1989.
5. Abrams HL. The relevance of Paleolithic diet in determining contemporary nutritional needs. *J Appl Nutr.* 1979;31:43-59.
6. Eaton SB, Konner MJ. Paleolithic nutrition: a consideration of its nature and current implications. *N Engl J Med.* 1985;312:283-289.
7. Truswell AS. Diet and nutrition of hunter-gatherers. In: *Health and Disease in Tribal Societies.* New York: Elsevier; 1977.
8. Ridker PM. On evolutionary biology, inflammation, infection, and the causes of atherosclerosis. *Circulation.* 2002;105:2-4.
9. Eaton SB. Humans, lipids and evolution. *Lipids.* 1992;27:814-820.
10. Eaton SB, Eaton SB III. Paleolithic vs. modern diets: selected pathophysiological implications. *Eur J Nutr.* 2000;39:67-70.
11. Cordain L, Miller JB, Eaton SB, Mann N, Holt SH, Speth JD. Plant-animal subsistence ratios and macronutrient energy estimations in worldwide hunter-gatherer diets. *Am J Clin Nutr.* 2000;71:682-692.
12. Cordain L, Eaton SB, Brand Miller J, Mann N, Hill K. The paradoxical nature of hunter-gatherer diets: meat-based, yet non-atherogenic. *Eur J Clin Nutr.* 2002;56 (suppl 1):S1-S11.
13. United States Department of Agriculture. The Food Guide Pyramid. Center for Nutrition Policy and Promotion. Home and Garden Bulletin 252. Washington, D.C., 2000. [http://www.pueblo.gsa.gov/cic\\_text/food/food-pyramid/main.htm](http://www.pueblo.gsa.gov/cic_text/food/food-pyramid/main.htm).
14. Cordain L. Cereal grains: humanity's double-edged sword. *World Rev Nutr Diet.* 1999;84:19-73.
15. Eaton SB, Nelson DA. Calcium in evolutionary perspective. *Am J Clin Nutr.* 1991;54(1 suppl):281S-287S.
16. Zohary D, Hopf M. Domestication of pulses in the Old World. *Science.* 1973;182:887-894.

17. Kurtzweil P. Nutritional info available for raw fruits, vegetables, fish. FDA Consumer Magazine. May, 1993. United States Health and Human Services, Food and Drug Administration, Rockville MD. <http://www.fda.gov/fdac/special/foodlabel/raw.html>.
18. Holt SA, Brand Miller JC, Petocz P. An insulin index of foods: the insulin demand generated by 1000-kJ portions of common foods. *Am J Clin Nutr*. 1997;66:1264-1276.
19. Thorburn AW, Brand JC, Truswell AS. Slowly digested and absorbed carbohydrate in traditional bushfoods: a protective factor against disease. *Am J Clin Nutr*. 1987;45:98-106.
20. United States Department of Agriculture, Economic Research Service. America's Eating Habits: Changes and Consequences. Agriculture Information Bulletin No. 750, Elizabeth Frazaio, ed., Washington, D.C., 1999. <http://www.ers.usda.gov/publications/aib750/aib750app.pdf>.
21. Cordain L, Watkins BA, Florant GL, Kehler M, Rogers L, Li Y. Fatty acid analysis of wild ruminant tissues: evolutionary implications for reducing diet-related chronic disease. *Eur J Clin Nutr*. 2002;56:1-11.
22. Heaney RP. Excess dietary protein may not adversely affect bone. *J Nutr*. 1998;128:1054-1057.
23. U.S. Department of Agriculture, Agricultural Research Service. 1997. Data tables: Results from USDA's 1994-96 Continuing Survey of Food Intakes by Individuals and 1994-96 Diet and Health Knowledge Survey, [Online]. ARS Food Surveys Research Group. Available (under "Releases"): <http://www.barc.usda.gov/bhnrc/foodsurvey/home.htm>.
24. Bridges PS. Skeletal biology and behavior in ancient humans. *Ev Anthropol*. 1995;4:112-120.
25. Ruff C, Trinklaus E, Walker A, Larsen CS. Postcranial robusticity in *Homo*. I: Temporal trends and mechanical interpretation. *Am J Phys Anthropol*. 1993;91:21-53.
26. Kricun ME. Edward B D Neuhauser Lecture. Paleoradiology of the prehistoric Australian aborigines. *AJR Am J Roentgenol*. 1994;163:241-247.
27. Barzel US. The skeleton as an ion exchange system: implications for the role of acid-base imbalance in the genesis of osteoporosis. *J Bone Min Res*. 1995;10:1431-1436.
28. Lutz J. Calcium balance and acid-base status of women as affected by increased protein intake and by sodium bicarbonate ingestion. *Am J Clin Nutr*. 1984; 39:281-288.
29. Sebastian A, Harris ST, Ottaway JH, Todd KM, Morris RC. Improved mineral balance and skeletal metabolism in postmenopausal women treated with potassium bicarbonate. *N Engl J Med*. 1994;33:1776-1781.
30. Remer T, Manz F. Potential renal acid load of foods and its influence on urine pH. *J Am Diet Assoc*. 1995;95:791-797.
31. Appel LJ, Moore TJ, Obarzanek E, Vollmer WM *et al*. A clinical trial of the effects of dietary patterns on blood pressure. *N Engl J Med*. 1997;336:1117-1124.
32. McNamara DJ, Kolb R, Parker TS, Batwin H, Samuel P, Brown CD, Ahrens EH. Heterogeneity of cholesterol homeostasis in man. Response to changes in dietary fat quality and cholesterol quantity. *J Clin Invest*. 1987;79:1729-1739.
33. Howell WH, McNamara DJ, Tosca MA, Smith BT, Gaines JA. Plasma lipid and lipoprotein responses to dietary fat and cholesterol: a meta-analysis. *Am J Clin Nutr*. 1997;65:1747-1764.
34. Schonfeld G, Patsch W, Rudel LL, Nelson C, Epstein M, Olson RE. Effects of dietary cholesterol and fatty acids on plasma lipoproteins. *J Clin Invest*. 1982;69:1072-1080.
35. Anonymous. Effects of dietary protein restriction on the progression of moderate renal disease in the Modification of Diet in Renal Disease Study. *J Am Soc Nephrol*. 1996;12:2616-2626.
36. Skov AR, Toubro S, Bulow J, Krabbe K, Parving HH, Astrup A. Changes in renal function during weight loss induced by high vs low-protein low-fat diets in overweight subjects. *Int J Obes Relat Metab Disord*. 1999;23:1170-1177.
37. Crovetti R, Porrini M, Santangelo A, Testolin G. The influence of thermic effect of food on satiety. *Eur J Clin Nutr*. 1998;52:482-488.
38. Stubbs RJ. Nutrition Society Medal Lecture. Appetite, feeding behaviour and energy balance in human subjects. *Proc Nutr Soc*. 1998;57:341-356.
39. Skov AR, Toubro S, Ronn B, Holm L, Astrup A. Randomized trial on protein vs carbohydrate in ad libitum fat reduced diet for the treatment of obesity. *Int J Obes Relat Metab Disord*. 1999;23:528-536.
40. Baba NH, Sawaya S, Torbay N, Habbal Z, Azar S, Hashim SA. High protein vs high carbohydrate hypoenergetic diet for the treatment of obese hyperinsulinemic subjects. *Int J Obes Relat Metab Disord*. 1999;23:1202-1206.
41. O'Dea K. Marked improvement in carbohydrate and lipid metabolism in diabetic Australian Aborigines after temporary reversion to traditional lifestyle. *Diabetes*. 1984;33:596-603.
42. O'Dea K, Traianedes K, Ireland P, Niall M, Sadler J, Hopper J, DeLuise M. The effects of diet differing in fat, carbohydrate, and fiber on carbohydrate and lipid metabolism in type II diabetes. *J Am Diet Assoc*. 1989; 89:1076-1086.
43. Wolfe BM, Giovannetti PM. Short term effects of substituting protein for carbohydrate in the diets of moderately hypercholesterolemic human subjects. *Metabolism*. 1991;40:338-343.
44. Wolfe BM, Giovannetti PM. High protein diet complements resin therapy of familial hypercholesterolemia. *Clin Invest Med*. 1992;15:349-359.
45. Wolfe BM, Piche LA. Replacement of carbohydrate by protein in a conventional-fat diet reduces cholesterol and triglyceride concentrations in healthy normolipidemic subjects. *Clin Invest Med*. 1999;22:140-148.
46. Seino Y, Seino S, Ikeda M, Matsukura S, Imura H. Beneficial effects of high protein diet in treatment of mild diabetes. *Hum Nutr Appl Nutr*. 1983;37A(3):226-230.
47. Piatti PM, Monti F, Fermo I, Baruffaldi L, Nasser R, Santambrogio G, Librenti MC, Galli-Kienle M, Pontiroli AE,



- Pozza G. Hypocaloric high-protein diet improves glucose oxidation and spares lean body mass: comparison to hypocaloric high-carbohydrate diet. *Metabolism*. 1994;43:1481-1487
48. Hu FB, Stampfer MJ, Manson JE, Rimm E, Colditz GA, Speizer FE, Hennekens CH, Willett WC. Dietary protein and risk of ischemic heart disease in women. *Am J Clin Nutr*. 1999;70:221-227.
  49. Stolzenberg-Solomon RZ, Miller ER III, Maguire MG, Selhub J, Appel LJ. Association of dietary protein intake and coffee consumption with serum homocysteine concentrations in an older population. *Am J Clin Nutr*. 1999; 69:467-475.
  50. Mann NJ, Li D, Sinclair AJ, Dudman NP, Guo XW, Elsworth GR, Wilson AK, Kelly FD. The effect of diet on plasma homocysteine concentrations in healthy male subjects. *Eur J Clin Nutr*. 1999;53:895-899.
  51. Mezzano D, Munoz X, Martinez C, Cuevas A, Panes O, Aranda E, Guasch V, Strobel P, Munoz B, Rodriguez S, Pereira J, Leighton F. Vegetarians and cardiovascular risk factors: hemostasis, inflammatory markers and plasma homocysteine. *Thromb Haemost*. 1999;81:913-917.
  52. Obarzanek E, Velletri PA, Cutler JA. Dietary protein and blood pressure. *JAMA*. 1996;275:1598-1603.
  53. Burke V, Hodgson JM, Beilin LJ, Giangiulioi N, Rogers P, Puddey IB. Dietary protein and soluble fiber reduce ambulatory blood pressure in treated hypertensives. *Hypertension*. 2001;38:821-826.
  54. Klag MJ, Whelton PK. The decline in stroke mortality. An epidemiologic perspective. *Ann Epidemiol*. 1993;3:571-575.
  55. Kinjo Y, Beral V, Akiba S, Key T, Mizuno S, Appleby P, Yamaguchi N, Watanabe S, Doll R. Possible protective effect of milk, meat and fish for cerebrovascular disease mortality in Japan. *J Epidemiol*. 1999;9:268-274.
  56. Krauss RM, Deckelbaum RJ, Ernst N, *et al*. Dietary guidelines for healthy American adults. *Circulation*. 1996;94:1795-1800.
  57. Denke MA, Breslow JL. Effects of a low fat diet with water and without intermittent saturated fat and cholesterol ingestion on plasma lipid, lipoprotein, and apolipoprotein levels in normal volunteers. *J Lipid Res*. 1988;29: 963-969.
  58. Dreon DM, Fernstrom HA, Miller B, Krauss RM. Apolipoprotein E isoform phenotype and LDL subclass response to a reduced-fat diet. *Arterioscler Thromb Vasc Biol*. 1995;15:105-111.
  59. Jeppesen J, Schaaf P, Jones C, Zhou M-Y, Ida Chen Y-D, Reaven GM. Effects of low-fat, high carbohydrate diets on risk factors for ischemic heart disease in postmenopausal women. *Am J Clin Nutr*. 1997;65:1027-1033.
  60. Mensink RP, Katan MB. Effect of dietary fatty acids on serum lipids and lipoproteins: a meta-analysis of 27 trials. *Arterioscler Thromb*. 1992;12:911-919.
  61. Grundy SM. Hypertriglyceridemia, insulin resistance, and the metabolic syndrome. *Am J Cardiol*. 1999;83:25F-29F.
  62. Foster-Powell K, Brand Miller J. International table of glycemic index. *Am J Clin Nutr*. 1995;62:871S-893S.
  63. Bjorck I, Liljeberg H, Ostman E. Low glycaemic-index foods. *B J Nutr*. 2000;83(suppl 1):S149-S155.
  64. Ebbeling CB, Ludwig DS. Treating obesity in youth: should dietary glycemic load be a consideration? *Adv Pediatr*. 2001;48:179-212.
  65. Salmeron J, Ascherio A, Rimm EB, Colditz GA, *et al*. Dietary fiber, glycemic load, and risk of NIDDM in men. *Diabetes Care*. 1997;20:545-550.
  66. Salmeron J, Manson JE, Stampfer MJ, Colditz GA, Wing AL, Willett WC. Dietary fiber, glycemic load, and risk of non-insulin-dependent diabetes mellitus in women. *JAMA*. 1997;277:472-477.
  67. Liu S, Willett WC, Stampfer MJ, Hu FB, Franz M, Sampson L, Hennekens CH, Manson JE. A prospective study of dietary glycemic load, carbohydrate intake, and risk of coronary heart disease in US women. *Am J Clin Nutr*. 2000;71: 1455-1461.
  68. Anderson JW, Smith BM, Gustafson NJ. Health benefits and practical aspects of high-fiber diets. *Am J Clin Nutr* 1994;59:1242S-1247S.
  69. Connor SL, Connor WE. Are fish oils beneficial in the prevention and treatment of coronary artery disease? *Am J Clin Nutr*. 1997;66:1020S-1031S.
  70. Gardner CD, Kraemer HC. Monounsaturated versus polyunsaturated dietary fat and serum lipids: a meta-analysis. *Arterioscler Thromb Vasc Biol*. 1995;15:1917-1927.
  71. Oliver MF. It is more important to increase the intake of unsaturated fats than to decrease the intake of saturated fats: evidence from clinical trials relating to ischemic heart disease. *Am J Clin Nutr*. 1997; 66:980S-986S.
  72. Nelson GJ, Schmidt PC, Kelley DS. Low-fat diets do not lower plasma cholesterol levels in healthy men compared to high-fat diets with similar fatty acid composition at constant caloric intake. *Lipids*. 1995;30: 969-976.
  73. Grundy SM. What is the desirable ratio of saturated, polyunsaturated, and monounsaturated fatty acids in the diet? *Am J Clin Nutr*. 1997;66:988S-990S.
  74. Yu S, Deer J, Etherton T, Kris-Etherton P. Plasma cholesterol-predictive equations demonstrated that stearic acid is neutral and monounsaturated fatty acids are hypocholesterolemic. *Am J Clin Nutr*. 1995;61:1129-1139.
  75. Garg A, Bonanome A, Grundy SM, Zhang Z-J, Unger RH. Comparison of a high- carbohydrate diet with a high-monounsaturated fat diet in patients with non-insulin dependent diabetes mellitus. *N Eng J Med*. 1988;319:829-834.
  76. National Cholesterol Education Program. Second Report of the Expert panel on Detection, Evaluation and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel II). *Circulation*. 1994;89:1333-1445.
  77. O'Byrne DJ, O'Keefe SF, Shireman RB. Low-fat, monounsaturated-rich diets reduce susceptibility of low density lipoproteins to peroxidation *ex vivo*. *Lipids*. 1998;33:149-157.

78. U.S. Department of Agriculture, Agricultural Research Service. 1997. Data tables: Intakes of 19 Individual Fatty Acids: Results from USDA's 1994-96 Continuing Survey of Food Intakes by Individuals, [Online]. ARS Food Surveys Research Group. Available (under "Releases"): <http://www.barc.usda.gov/bhnrc/foodsurvey/home.htm>.
79. de Lorgeril M, Salen P, Martin JL, Monjaud I, Delaye J, Mamelle N. Mediterranean diet, traditional risk factors, and the rate of cardiovascular complications after myocardial infarction: final report of the Lyon Diet Heart Study. *Circulation*. 1999;99:779-785.
80. Singh RB, Niaz MA, Sharma JP, Kumar R, Rastogi V, Moshiri M. Randomized, double-blind, placebo-controlled trial of fish oil and mustard oil in patients with suspected acute myocardial infarction: the Indian experiment of infarct survival-4. *Cardiovasc Drugs Ther*. 1997;11:485-491.
81. von Schacky C, Angerer P, Kothny W, Theisen K, Mudra H. The effect of dietary omega-3 fatty acids on coronary atherosclerosis: a randomized, double-blind, placebo-controlled trial. *Ann Intern Med*. 1999;130:554-562.
82. GISSI-Prevention Investigators. Dietary supplementation with n-3 polyunsaturated fatty acids and vitamin E after myocardial infarction: results of the GISSI-Prevenzione trial. Gruppo Italiano per lo Studio della Sopravvivenza nell'Infarto miocardico. *Lancet*. 1999;354:447-455.
83. Antonios TF, MacGregor GA. Salt—more adverse effects. *Lancet*. 1996;348:250-251.
84. Massey LK, Whiting SJ. Dietary salt, urinary calcium, and kidney stone risk. *Nutr Rev*. 1995;53:131-139.
85. Boushey CJ, Beresford SA, Omenn GS, *et al*. A quantitative assessment of plasma homocysteine as a risk factor for vascular disease: probable benefits of increasing folic acid intakes. *JAMA*. 1995;274:1049-1057.
86. Robinson K, Arheart K, Refsum H, *et al*. Low circulating folate and vitamin B6 concentrations: risk factors for stroke, peripheral vascular disease, and coronary artery disease: European COMAC Group. *Circulation*. 1998;97:437-443.

**Latest 21st Century Medical Advances  
in the Diagnosis & Treatment of  
Fibromyalgia,  
Chronic Fatigue Syndrome  
and Related Illnesses**

**Cutting-edge Conference for Physicians and Healthcare Professionals on Clinical and Office-Based Applications**

**September 19 - 21, 2002**

**Radisson Hotel - LAX  
Los Angeles, CA**

Co-sponsored by National Fibromyalgia Association,  
American Association for Chronic Fatigue Syndrome,  
Foundation for Care Management &  
The Healthy Foundation

For more information: [www.AdMedCon.com](http://www.AdMedCon.com)  
or call 800-863-5085



**Educational Curriculum Includes:**

- Metabolic Treatments of CFIDS – Dr. Jacob Teitelbaum
- New Discoveries in Treating CFIDS – Dr. Paul Cheney
- Diagnosing and Treating Myofascial Pain – Dr. Hal Blatman
- Immune Therapies – Dr. Michael Rosenbaum
- Psychological Issues – Dr. Hyla Cass
- Treatment Strategies for Post-Partum CFIDS – Dr. Dean Raffelock
- Laboratory Markers for CFIDS – Dr. Ari Vojdani
- Reimbursement Issues for Patients – Scott Davis, Attorney
- Anti-Virals for CFIDS – Dr. Sbari Lieberman
- Importance of Calcium and Magnesium – Dr. Nan Fuchs
- Practice Management Workshop – Dr. Jacob Teitelbaum
- Clinical Outcome Study & Research for FM – Dr. Robert Bradford
- Enhancing Biological Methylation to Mitigate the Symptoms of Fibromyalgia – Dr. Todd Ovokaitys
- Clinical Outcome Study & Research for FM – Dr. Serafina Corsello